

Multi-scale analysis of hermatypic coral assemblages at Mexican Central Pacific

Joicye Hernández-Zulueta, Fabián A. Rodríguez-Zaragoza, Rubén Araya,
Ofelia Vargas-Ponce, Alma P. Rodríguez-Troncoso, Amílcar L. Cupul-Magaña,
Leopoldo Díaz-Pérez, Eduardo Ríos-Jara, Marco Ortiz

Supplementary material




Fig. S1. – Latitude of the different zones of the Mexican Central Pacific with their coral species richness. Codes by zones: IIN (Isla Isabel); IMN (Islas Marietas); BCH (Bahía Chamela); CUM (Bahía Cuastecomate-Punta Melaque); BCE (Bahía Ceníceros); FB (Faro de Bucerías) and CAC (Caleta de Campos).

Table S1. – Richness and coverages (%) of hermatypic corals species in the different sampling sites of the Mexican Central Pacific. Codes by states: NAY (Nayarit); JAL (Jalisco); COL (Colima) and MICH (Michoacán). Codes by Zones: IIN (Isla Isabel); IMN (Islas Marietas); BCH (Bahía Chamela); CUM (Bahía Cuastecomate-Punta Melaque); BCE (Bahía Ceníceros); FB (Faro de Bucerías) and CAC (Caleta de Campos). Codes by sites: PRO (Punta Rocosa); CSN (Caleta Sin Nombre); LMI (Las Monas Interno); LME (Las Monas Externo); CFS (Costa Fragata Somero); CFP (Costa Fragata Profundo); BRA (Bahía Rabijuncos); EPE (Ensenada de Pescadores); POC (Pocitas); PA (Playa del Amor); PP (Plataforma Pavonas); TAM (Túnel Amarradero); ZR (Zona de Restauración); ZRS (Zona de Restauración Sur); CDM (Cueva del Muerto); P (Pelicanos); Lpj (La Pajarera); Lpa (La Palma); EEL (El Elefante); EC (Ensenada Cocinas); IC (Isla Cocinas); TE (Tenacatita); LV (La Virgencita); CU (Cuastecomatito); LMO (Las Monjas 1); LMT (Las Monjas 2); CRZ (Carrizales); PM (Paraíso Montemar); LR (L'Recif); EE (Elefante); PB (Punto B); LB (La Boquita); IPE (Isla Pájaros Este); BH (Barco Hundido); IPO (Isla Pájaro Oeste); IP (Isla Protegida); ER (El Relis); EF (El Faro); LE (La Ensenada); ET (El Teolan) and MC (Manto Carrizo). Codes by coral species: *Pc*, *P. capitata*; *Pd*, *P. damicornis*; *Pef*, *P. effusus*; *Pey*, *P. eydouxi*; *Pi*, *P. inflata*; *Pm*, *P. meandrina*; *Pv*, *P. verrucosa*; *Pl*, *P. lobata*; *Pp*, *P. panamensis*; *Pcl*, *P. clavus*; *Pdu*, *P. duerdeni*; *Pg*, *P. gigantea*; *Pva*, *P. varians*; *Pst*, *P. stellata* and *Psu*, *P. superficialis*. S, Richness of coral species.

ST	ZN	SI	<i>Pc</i>	<i>Pd</i>	<i>Pef</i>	<i>Pey</i>	<i>Pi</i>	<i>Pm</i>	<i>Pv</i>	<i>Pl</i>	<i>Pp</i>	<i>Pcl</i>	<i>Pdu</i>	<i>Pg</i>	<i>Pva</i>	<i>Pst</i>	<i>Psu</i>	S		
NAY	IIN	PRO	0.5		0.6				3.2	3.9		2.6				0.8	0.3	7		
		CSN		0.4					5.0	6.1									2	
		LMI	1.7	0.5	0.1				4.8	1.0				0.2				0.1	7	
		LME		0.2					2.3	0.2									3	
		CFS	9.0	6.1					5.4	1.4									4	
		CFP							0.2	0.1									2	
		BRA		0.1				0.1	3.2	8.0	0.1				0.1		0.2		7	
		EPE	4.1	2.4			0.2		12.3	4.0							0.1	0.5	7	
		POC							0.9	2.8							0.1	0.4	4	
		PA							0.1	4.6		2.8							3	
		PP								2.0		1.8		12.3					3	
		TAM								1.4		1.2		3.1					3	
		ZR	0.2	0.1			0.5		0.3	6.1		3.0							6	
		ZRS	3.4	3.4					0.1	15.7		3.2		0.5					6	
JAL	BCH	CDM	0.2	1.4					8.9		2.9							4		
		P	1.5	8.5		1.7			12.5									4		
		LPj	0.7	0.4		1.3				4.4		1.7							5	
		Lpa	21.6	6.4				0.2	20.7	0.2	0.7								6	
		EEL							1.6	0.7	1.1								3	
		EC		0.3				0.2	0.9	0.4	0.8								5	
		IC	1.2						0.1		1.7			0.9					4	
		TE	16.9	40.8					8.6	1.2	0.3	0.2							6	
		LV	1.5	9.9					0.6	9.7	0.9	0.2							6	
		CU	5.4	36.5					1.3	2.5	0.6	0.9		0.5					7	
		LMO	7.4	41.2					1.9	2.6	0.1	0.1		0.1					7	
		LMT	0.3	1.4							0.5	0.7							4	
		COL	BCE	CRZ	12.5	19.1		0.2	0.4	0.3	19.3	0.3	0.2		0.3	0.3	0.1			11
				PM		0.1				0.2		1.1	1.5							4
LR										3.0	1.0							2		
EE				0.1					0.1	0.1	0.4							4		
BSG	PB		0.1	7.5					3.2	0.3	0.2			0.3					6	
	LB		0.2	9.0					1.3	4.8	0.3			0.1					5	
	IPE		2.5	0.5						0.2		9.3							4	
	BH											3.3							1	
MICH	FB	IPO									8.5							1		
		IP									0.8							1		
		ER										1.2						1		
		EF										3.9						1		
		LE		0.3					0.1		2.5			0.8				4		
		ET								1.0		0.2						2		
		MC										1.3						1		

Table S2. – Results of the ANOVA and PERMANOVA pairwise test. Codes by the states: NAY (Nayarit); JAL (Jalisco); COL (Colima) and MICH (Michoacán). Codes by zones: IIN (Isla Isabel); IMN (Islas Marietas); BCH (Bahía Chamela); CUM (Bahía Cuastecomate-Punta Melaque); BCE (Bahía Ceníceros); FB (Faro de Bucerías) and CAC (Caleta de Campos). Bold numbers correspond to a statistical significance $p < 0.05$.

	Group	t	P(perm)		Group	t	P(perm)	
ANOVA								
Species richness (S)	MICH vs. COL	2.7209	0.025	Shannon diversity (H')	MICH vs. COL	8.4951	0.0002	
	MICH vs. JAL	6.5742	0.0001		MICH vs. JAL	15.238	0.0001	
	MICH vs. NAY	2.6789	0.0154		MICH vs. NAY	6.0826	0.0001	
	COL vs. JAL	1.4477	0.1761		COL vs. JAL	1.3134	0.2351	
	COL vs. NAY	0.3879	0.7139		COL vs. NAY	0.5467	0.6015	
	JAL vs. NAY	1.9694	0.062		JAL vs. NAY	1.601	0.1281	
Live coral cover (LCC)	MICH vs. COL	1.3428	0.0207					
	MICH vs. JAL	3.4174	0.0036					
	MICH vs. NAY	2.3125	0.0335					
	COL vs. JAL	1.1252	0.2823					
	COL vs. NAY	0.0642	0.9505					
	JAL vs. NAY	1.8369	0.0789					
Coral assemblages by state	MICH vs. COL	3.1900	0.0009	Coral assemblages by zone	IIN vs. IMN	3.8892	0.0003	
	MICH vs. JAL	4.4237	0.0003		BCH vs. CUM	1.7266	0.0630	
	MICH vs. NAY	5.1706	0.0001		BCE vs. SGO	0.7083	0.6822	
	COL vs. JAL	1.3824	0.1461		FB vs. CAC	1.0539	0.3550	
	COL vs. NAY	2.1246	0.0052					
	JAL vs. NAY	2.2036	0.0053					
Benthonic habitat structure by state	MICH vs. COL	1.5571	0.0670	Benthonic habitat structure by zone	IIN vs. IMN	1.1832	0.2462	
	MICH vs. JAL	1.6078	0.0537		BCH vs. CUM	1.7558	0.0034	
	MICH vs. NAY	2.5896	0.0001		BCE vs. SGO	0.7347	0.6972	
	COL vs. JAL	1.1812	0.2477		FB vs. CAC	1.4774	0.1547	
	COL vs. NAY	2.5816	0.0007					
	JAL vs. NAY	2.3814	0.0011					

Table S3. – Results of the similarity percentage analysis (SIMPER) and coverages (%) of coral species by state. For SIMPER, only those species contributing 90% of the average dissimilarity (Ave. Dis.) of coral species between states of the Mexican Central Pacific are presented. AC, specie cover per state; C, percentage of contribution to the average dissimilarity; CC, percentage of accumulated contribution to the average dissimilarity. Codes: NAY (Nayarit); JAL (Jalisco); COL (Colima); and MICH (Michoacán).

	AC	AC	C	CC		AC	AC	C	CC
Ave. Dis. = 76.83%	MICH	NAY			Ave. Dis. = 57.20%	COL	JAL		
<i>P. verrucosa</i>	0.11	1.23	31.53	31.53	<i>P. damicornis</i>	1.23	1.18	22.04	22.04
<i>P. panamensis</i>	1.21	0.51	25.74	57.27	<i>P. verrucosa</i>	0.86	1.09	19.05	41.09
<i>P. lobata</i>	0	0.54	14.21	71.49	<i>P. capitata</i>	0.54	0.85	15.88	56.98
<i>P. capitata</i>	0.07	0.37	7.74	79.23	<i>P. lobata</i>	0.83	0.34	15.32	72.3
<i>P. damicornis</i>	0.08	0.33	7.62	86.85	<i>P. panamensis</i>	0.59	0.7	10.53	82.83
<i>P. gigantea</i>	0.05	0.16	5.25	92.1	<i>P. meandrina</i>	0.12	0.43	7.97	90.81

Table S4. – Results of the similarity percentage analysis (SIMPER) and coverages (%) of coral species per zone. For SIMPER, only those species contributing 90% of the average dissimilarity (Ave. Dis.) of coral species between zones of the Mexican Central Pacific are presented. AC, percentage of average cover per zone; C, percentage of contribution to the average dissimilarity; CC, percentage of accumulated contribution to the average dissimilarity. Codes: IIN (Isla Isabel); IMN (Islas Marietas); CAC (Caleta de Campos) and FB (Faro de Bucerías).

	AC	AC	C	CC		AC	AC	C	CC
Ave. Dis. = 71.03%	IMN	IIN			Ave. Dis. = 29.52%	CAC	FB		
<i>P. panamensis</i>	1.18	0.01	26.41	26.41	<i>P. panamensis</i>	1.04	1.43	60.04	60.04
<i>P. lobata</i>	0	0.94	19.17	45.58	<i>P. verrucosa</i>	0.14	0.08	17.38	77.42
<i>P. verrucosa</i>	1.28	1.2	15.88	61.45	<i>P. damicornis</i>	0.07	0.1	8.98	86.4
<i>P. capitata</i>	0.27	0.44	9.61	71.06	<i>P. capitata</i>	0	0.15	8.46	94.87
<i>P. damicornis</i>	0.28	0.37	9.16	80.22					
<i>P. gigantea</i>	0.33	0.04	7.74	87.96					
<i>P. stellata</i>	0	0.18	3.19	91.15					

Table S5. – Results of the similarity percentage analysis (SIMPER) and coverages (%) of coral species per site. For SIMPER, only those species contributing 90% of the average dissimilarity (Ave. Dis.) of coral species between sampling sites of the Mexican Central Pacific are presented. AC, percentage of average cover per site; C, percentage of contribution to the average dissimilarity; CC, percentage of accumulated contribution to the average dissimilarity. Codes: CFS (Costa Fragata Somero); CFP (Costa Fragata Profundo); CSN (Caleta Sin Nombre); POC (Pocitas); ZRS (Zona de Restauración Sur); TAM (Túnel Amarradero); ZR (Zona de Restauración); PA (Playa del Amor); P (Pelícanos); IC (Isla Cocinas); EEL (El Elefante); EC (Ensenada Cocinas); LMT (Las Monjas 2); TE (Tenacatita); CU (Cuastecomatito); LMO (Las Monjas 1); CRZ (Carrizales); LR (L Recif); PM (Paraíso Montemar); EE (Elefante); LB (La Boquita); PB (Punto B); ET (El Teolan); LE (La Ensenada); ER (El Relis); MC (Manto Carrizo); IPO (Isla Pájaro Oeste); IP (Isla Protegida); and BH (Barco Hundido).

	AC	AC	C	CC		AC	AC	C	CC
Isla Isabel									
Ave. Dis. = 77.49%	CFS	CFP			Ave. Dis. = 33.29%	CSN	POC		
<i>P. capitata</i>	1.69	0	38.32	38.32	<i>P. verrucosa</i>	1.44	0.77	39.63	39.63
<i>P. damicornis</i>	1.41	0	32.53	70.85	<i>P. lobata</i>	1.54	1.26	23.49	63.12
<i>P. verrucosa</i>	1.26	0.53	16.1	86.94	<i>P. superficialis</i>	0	0.29	12.94	76.06
<i>P. lobata</i>	0.54	0.3	13.06	100	<i>P. damicornis</i>	0.3	0	12.79	88.85
					<i>P. stellata</i>	0	0.23	11.15	100
Islas Marietas									
Ave. Dis= 60.61%	ZRS	TEMB			Ave. Dis.=25.92%	ZR	PA		
<i>P. verrucosa</i>	1.98	0.46	34.83	34.83	<i>P. panamensis</i>	1.28	1.13	22.88	22.88
<i>P. capitata</i>	1.12	0	24.69	59.52	<i>P. eydouxi</i>	0.41	0	21.56	44.44
<i>P. damicornis</i>	0.83	0	16.23	75.74	<i>P. verrucosa</i>	1.55	1.36	20.24	64.68
<i>P. gigantea</i>	0.25	0.53	13.14	88.89	<i>P. meandrina</i>	0.22	0.13	15.94	80.62
<i>P. panamensis</i>	1.3	0.97	7.98	96.86	<i>P. damicornis</i>	0.16	0	9.85	90.47
Bahía Chamela									
Ave. Dis= 88.55%	P	IC			Ave. Dis= 23.81%	EEL	EC		
<i>P. damicornis</i>	1.47	0	24.7	24.7	<i>P. lobata</i>	0.89	0.59	27.49	27.49
<i>P. verrucosa</i>	1.72	0.22	24.36	49.06	<i>P. meandrina</i>	0	0.29	21.91	49.41
<i>P. panamensis</i>	0	1.11	18.99	68.05	<i>P. damicornis</i>	0	0.33	21.3	70.7
<i>P. gigantea</i>	0	0.72	13.24	81.29	<i>P. verrucosa</i>	1	0.97	18.15	88.86
<i>P. capitata</i>	0.74	0.46	11.27	92.56	<i>P. panamensis</i>	1	0.94	11.14	100
Bahía Cuastecomate- Punta Melaque									
Ave. Dis= 71.43%	LMT	TE			Ave. Dis= 20.52%	CU	LMO		
<i>P. damicornis</i>	0.78	2.45	25.27	25.27	<i>P. panamensis</i>	0.96	0.22	24.46	24.46
<i>P. capitata</i>	0.33	1.95	23.95	49.22	<i>P. meandrina</i>	0.75	1.12	19.17	43.62
<i>P. meandrina</i>	0	1.65	23.43	72.66	<i>P. damicornis</i>	2.44	2.36	17	60.63
<i>P. verrucosa</i>	0	0.75	10.37	83.02	<i>P. lobata</i>	0.39	0.22	14.53	75.15
<i>P. panamensis</i>	0.88	0.29	9.69	92.72	<i>P. duerdeni</i>	0.36	0.22	13.83	88.98
					<i>P. capitata</i>	1.52	1.59	7.61	96.59
Bahía Ceníceros									
Ave. Dis. = 87.72%	CRZ	LR			Ave. Dis. = 30.48%	PM	LR		
<i>P. verrucosa</i>	2.05	0	24.2	24.2	<i>P. panamensis</i>	1.07	0.74	33.63	33.63
<i>P. damicornis</i>	2.02	0	23.82	48.02	<i>P. damicornis</i>	0.45	0	29.3	62.93
<i>P. capitata</i>	1.75	0	20.11	68.12	<i>P. lobata</i>	1.01	1.24	19.57	82.5
<i>P. lobata</i>	0.44	1.24	9.73	77.86	<i>P. meandrina</i>	0.29	0	17.5	100
<i>P. panamensis</i>	0.3	0.74	7.55	85.41					
<i>P. meandrina</i>	0.33	0	3.85	89.26					
<i>P. inflata</i>	0.34	0	3.44	92.7					
Bahía Santiago									
Ave. Dis. = 57.88%	EE	LB			Ave. Dis. =38.48%	LB	PB		
<i>P. damicornis</i>	0.33	1.57	35.51	35.51	<i>P. lobata</i>	1.26	0.52	26.86	26.86
<i>P. lobata</i>	0.33	1.26	27.06	62.57	<i>P. verrucosa</i>	0.56	0.94	24.41	51.26
<i>P. verrucosa</i>	0.67	0.56	16.06	78.64	<i>P. damicornis</i>	1.57	1.63	14.07	65.33
<i>P. panamensis</i>	0.87	0.42	13.69	92.33	<i>P. panamensis</i>	0.42	0.67	13.33	78.67
					<i>P. gigantea</i>	0.11	0.33	10.76	89.43
					<i>P. capitata</i>	0.18	0.27	10.57	100
Faro de Bucerías									
Ave. Dis. = 61.70%	ET	LE			Ave. Dis. = 13.87%	ER	MC		
<i>P. panamensis</i>	0.44	1.25	42.56	42.56	<i>P. panamensis</i>	0.98	0.99	100	100
<i>P. verrucosa</i>	0.66	0.22	35.22	77.78					
<i>P. gigantea</i>	0	0.41	12.63	90.41					
Caleta de Campos									
Ave. Dis. = 35.92%	IPO	IP			Ave. Dis. = 11.16%	BH	IP		
<i>P. panamensis</i>	1.53	1.04	42.26	42.26	<i>P. panamensis</i>	1.32	1.04	100	100
<i>P. capitata</i>	0.55	0	26.43	68.68					
<i>P. damicornis</i>	0.36	0	17.31	85.99					
<i>P. verrucosa</i>	0.29	0	14.01	100					

Table S6. – Coverages (%) of the morphofunctional groups per site. Codes by sites: PRO (Punta Rocosa); CSN (Caleta Sin Nombre); LMI (Las Monas Interno); LME (Las Monas Externo); CFS (Costa Fragata Somero); CFP (Costa Fragata Profundo); BRA (Bahía Rabijuncos); EPE (Ensenada de Pescadores); POC (Pocitas); PA (Playa del Amor); PP (Plataforma Pavonas); TAM (Túnel Amarradero); ZR (Zona de Restauración); ZRS (Zona de Restauración Sur); CDM (Cueva del Muerto); P (Pelicanos); Lpj (La Pajarera); Lpa (La Palma); EEL (El Elefante); EC (Ensenada Cocinas); IC (Isla Cocinas); TE (Tenacatita); LV (La Virgencita); CU (Cuastecomatito); LMO (Las Monjas 1); LMT (Las Monjas 2); CRZ (Carrizales); PM (Paraíso Montemar); LR (L Recif); EE (Elefante); PB (Punto B); LB (La Boquita); IPE (Isla Pájaros Este); BH (Barco Hundido); IPO (Isla Pájaro Oeste); IP (Isla Protegida); ER (El Relis); EF (El Faro); LE (La Ensenada); ET (El Teolan); and MC (Manto Carrizo). Codes by morphofunctional groups: LCC, Live coral cover; DSC, Soft coral; HC, Hydrocoral; RSC, Sponge; TUC, Turf; ACA, Articulate algae; CCA, Crustose calcareous algae; FMA, Fleahy macroalgae; RU, Rubble; DCC, Death coral; RO, Rock; SS, Sand; BAC, Barnacles.

	LCC	DSC	HC	RSC	TUC	ACA	CCA	FMA	RU	DCC	RO	SS	BAC
PRO	11.9	0.6	1.4	1.0	50.9		7.1	0.5	0.8	0.1	16.9	8.8	
CSN	11.5		0.7	0.1	48.4		6.3	4.3	0.6		9.5	18.7	
LMI	8.4	0.4	2.5	1.7	48.7		1.8	0.3	0.2	0.3	30.4	5.2	
LME	2.7	0.1	3.1	1.7	44.3		2.4	0.3	0.6		29.9	14.9	
CFS	21.9		0.7	0.2	52.7		6.9	14.9	2.0	0.1		0.6	
CFP	0.3			0.5	58.3		0.1	2.5			25.0	13.2	
BRA	11.6	0.1	0.2	0.5	60.9	0.1	5.4	14.1	0.1		3.5	3.5	
EPE	23.5			0.1	39.5		0.6	0.4	2.5	0.5	19.1	13.8	
POC	4.2		0.4	0.7	59.4		1.3	0.3	0.1	0.1	16.2	17.2	
PA	7.5		0.7	0.5	18.8	0.4	6.4	0.1	0.4	0.3	61.0	3.9	
PP	16.0	0.5	3.2	4.6	50.6		5.9		3.0	0.8	3.9	11.5	
TAM	5.8	0.5	0.8	0.6	45.5		3.7	0.4	2.4	1.1	16.2	23.0	
ZR	10.2		0.4	0.2	18.6		7.7	7.0	1.2	0.4	48.4	5.7	
ZRS	26.4				20.1		3.5	8.0	4.7	0.2	35.2	2.0	
CDM	13.3		0.6	0.4	55.6		7.8	3.6	3.2	0.9	6.6	8.1	
P	24.2					0.1	0.2	0.8	23.6	11.5	33.4	6.2	
Lpj	8.5	0.3				1.2	0.7	12.8		0.1	74.1	2.5	
Lpa	49.9		0.1	0.7	14.3	2.1	2.7	15.6	4.7		7.3	2.5	
EEL	3.5	0.5	0.6	1.4	33.7	4.8	16.0	9.3	2.5		13.5	14.3	
EC	2.7	0.7	2.2	1.9	46.9	0.7	17.7	4.1	0.9		2.8	0.7	18.9
IC	3.9	0.7				0.1		0.4	0.0		76.8	18.1	
TE	67.9				7.1	0.9	0.7	7.5	10.9		0.0	4.9	
LV	22.9				24.3	3.0	10.8	33.1	2.0		0.8	3.1	
CU	47.7			0.1	35.2	2.8	3.1	7.0	2.2		0.5	1.3	
LMO	53.3			0.1	19.3	1.8	8.0	13.9	1.5		1.2	0.9	
LMT	2.9	0.1	1.3	1.7	59.4		17.2	2.6	1.4		7.1	6.3	
CRZ	53.0		0.2		10.4	2.1	3.3	25.3	1.6		0.1	3.9	
PM	2.9	0.1	0.8	2.7	59.1	10.9	11.0	1.9	1.7		4.3	4.7	
LR	4.0	0.3	1.6	1.3	57.9	0.4	26.2				3.6	4.7	
EE	0.7	0.1	2.0	1.1	18.3	7.6	1.4	57.0	2.9		3.6	5.5	
PB	11.6		0.3	4.3	39.9	9.1	7.7	14.2	7.5		2.1	3.3	
LB	15.7			1.6	9.4	1.1	0.4	61.7	1.4		1.4	7.3	
IPE	12.5				50.1	0.7	12.6	6.6	3.3		4.6	9.7	
BH	3.3	0.3			66.3		21.3	1.3			0.7	6.9	
IPO	8.5	0.1	0.5		23.0	2.4	14.5	9.3	2.5		25.6	13.6	
IP	0.8	1.5	0.1	1.5	62.2		7.3	3.0			3.3	20.2	
ER	1.2	4.4	1.1	0.4	19.0	0.2	3.0	14.0	0.3		41.5	14.9	
EF	3.9				34.7	0.9	15.5	35.4	0.1		5.1	4.4	
LE	3.7	0.1		0.1		3.5	1.5	61.7			24.7	4.7	
ET	1.2	3.9	0.5	1.9	56.3		12.0	6.3			8.5	9.3	
MC	1.3			0.5	21.9	3.2	10.5	38.1			1.2	23.3	